



Insect and Mite Galls:

Myths and Misconceptions

In Part II of this series, we learn about insect and mite galls, including the Six Laws of such galls.

By Joe Boggs and Jim Chatfield

In the first installment of this series on plant galls (May 2015), we talked about the difference between gall-like structures and true galls, including bacterial crown galls, fungal galls, leaf/petiole galls, flower/fruit galls, bud galls and stem galls. Galls galore! But there's more.

Let's dig deeper into insect and mite (arthropods) galls. Unlike bacterial crown galls, which are a mass of plant cells that have been modified by bacterial DNA, or fungal galls, which are an assemblage of fungal cells intertwined with plant cells, galls produced by insects and mites are constructed entirely of plant cells.

Dominant arthropod gall-makers include insects belonging to three orders: Hymenoptera (wasps, sawflies); Hemiptera (aphids/adelgids, phylloxerans, psyllids); and Diptera (midge flies). The only mites capable of inducing galls are eriophyids (order Trombidiformes, family Eriophyidae).

There are more than 2,000 species of insect gall-makers in the U.S.; however, three quarters belong to only two families: Cynipidae ("gall wasps") and Cecidomyiidae ("gall midges"). Note the "cecido-" in the name of the gall midge family; Cecidology is the scientific study of plant galls. Of the over 800 different gall-makers on oaks, more than 700 are gall wasps. In other words, when trying to identify an insect gall, keep in mind that there is a high probability it was produced by a gall wasp or gall midge.


Although insect and mite gall formation is not entirely understood, researchers theorize there are two possible pathways. Some gall researchers believe certain types of plant gall growth are directed by the feeding activity of the gall-maker. The galls are produced by a combination of constant but subtle feeding irritation, perhaps coupled with the release of chemical inducers by the gall-maker.

Certain eriophyid mites provide an example. These unusual mites are much smaller than spider mites (you need at least 40x magnification to see them); however, both types of mites use their

sharp, piercing mouthparts (chelicerae) to rupture plant cells so they can feed on the contents. Only the feeding activity of some species of eriophyid mites induces gall growth; there are no spider mite gall-makers. This gall-growth pathway may explain how simple felt-like erineum patches (a.k.a. "erineum galls") develop under the direction of a number eriophyid mite species. However, it does not explain how highly organized plant gall structures develop.

Some types of insect and mite galls are composed of complex plant structures and may include functional plant

Stem Gall Growth




Graphic by Joe Boggs, OSU Extension

Top of page, fuzzy, felt-like erineum patch galls appear on a birch leaf.

All photos: Courtesy of Joe Boggs



Mossy rose gall is a good example of a plurilocular, unilarval gall (multiple chambers but only one larva per chamber), while hickory phylloxera petiole gall is a typical unilocular, multilarval gall (one chamber hosting several larvae).



The large burl affecting this walnut is not a true gall. Below, the oak anthracnose fungus, *Apiognomonia quercina*, is seen infecting the plant tissue of the small oak apple gall..

organs such as nectaries. These types of plant galls require a different gall-growth theory; one that includes the ability of the gall-maker to turn plant genes on and off as they direct plant cells to form highly organized plant structures.

Research has revealed that some gall-making insects and mites produce chemical replicas of plant hormones, or “plant hormone analogs” meaning the molecules are nothing like plant hormones but the plant’s response is the same as with plant hormones. The gall-forming process is usually initiated by the female when she injects gall-inducing chemicals into the plant along with her eggs. The eggs themselves may ooze gall-inducing chemicals and once the eggs hatch, the interaction continues with the immature gall-makers continuing to exude chemicals to direct plant growth to suit their needs.

The resulting galls provide both a protective home and nourishment for the next generation of gall-maker. The continual direction of gall growth by the gall-maker using chemicals to turn plant genes on and off speaks to why some find insect and mite plant galls so fascinating.

The chemicals exuded by gall-makers can only act upon meristematic plant cells, such as the cambial cells mentioned in Part 1 of this series or the meristematic cells in leaf buds; the precursors to leaf cells. Under the influence of chemicals exuded by a gall-maker, the meristematic

cells that were originally destined to become leaf cells begin marching to a different drummer. Once the errant leaf cells fall under the chemical spell of a gall-maker, there is no turning back—they will become gall tissue. This means that gall formation cannot occur once meristematic leaf bud cells are committed to becoming leaf tissue; it’s one reason the leaf-gall season begins in the spring! However, once the galls start growing they will continue to grow, even after the leaves fully expand.

Stem galls that arise from cambial tissue present a different scenario. Since cambial cells remain “free agents” throughout the growing season, galls can be formed from these cells anytime during the growing season, although most stem galls also start growing early in the season to provide ample time for the gall-maker to complete its development.

Insect and mite gall identification

Identifying insect and mite galls is challenging because of the limited number of accurate and updated gall identification resources, both online and printed. This is particularly true for galls found in North America; the European gall literature is more robust.

The existing gall identification resources tend to follow the same general outline. Most begin by separating galls based on the gall-maker, such as wasp



galls, midge galls, and so on. While it is ultimately important to know the gall-maker, gall identification usually starts with the gall itself.

We’ll provide a quick recap from Part I: The first step is to consider where the gall is found on the plant. Most identification resources use the following locations: leaf/petiole galls; flower/fruit galls; bud galls; stems galls; and root galls.

Next, galls are described based on their structure such as general appearance (for example, ball-like, hairy, etc.), number of chambers, and the number of gall-makers housed in the chambers. Recall that *unilocular* galls have only one chamber; *plurilocular* galls have multiple chambers. *Unilarval* galls only have one gall-maker per chamber; *multilarval* galls have more than one gall-maker per chamber. The aforementioned ball-like larger oak apple gall is a unilocular, unilarval gall. The hairy-looking mossy rose

Continued on page 24

The continual direction of gall growth by the gall-maker using chemicals to turn plant genes on and off speaks to why some find insect and mite plant galls so fascinating.



Looking like a green apple, a large oak apple gall contains a wasp larva; once the larva pupates, the mature gall turns tannish brown and appears empty.

Continued from page 23

gall produced under the direction of the gall wasp, *Diplolepis rosae*; a species first described by Carl Linnaeus, is a plurilocular, unilarval gall with many chambers but only one larva per chamber.

The ball-like hickory petiole galls produced by several phylloxera species (*Phylloxera* spp.) are usually unilocular, multilarval galls; there is a single chamber housing many of these aphid relatives.

Insect and mite gall laws

While the highly organized plant galls produced under the direction of many insect and mite gall-makers represent a wide range of forms and locations, there are certain consistencies that can be summarized as “gall laws.”

The First Gall Law: Galls are abnormal plant growths produced under the direction of a living gall-maker; they do not arise spontaneously, nor are they in response to plant wounding that does not involve a gall-maker.

The First Gall Law: Galls are abnormal plant growths produced under the direction of a living gall-maker; they do not arise spontaneously, nor are they in response to plant wounding that does not involve a gall-maker. This law removes certain tree growth such as “burls” from the gall arena. It is believed these abnormal plant growths are the result of run-

away plant hormone production; they are not considered “true galls.”

The Second Gall Law: Insect and mite galls are abnormal plant structures that are composed entirely of plant tissue; they’re not part of the gall-maker. Infections by fungal plant pathogens can illustrate that insect and mite plant galls are indeed plant structures. The oak anthracnose fungus, *Apiognomonia quercina*, normally infects the leaf cells of its namesake host producing blackened, necrotic leaf tissue. The fungus can also infect the plant tissue of the small oak apple gall produced under the direction of the gall wasp, *Cynips clivorum*, because the gall is constructed from hijacked leaf cells. The resulting tissue necrosis makes the gall appear to slowly dissolve away.

The Third Gall Law: Galls can only be formed from meristematic plant tissue and once plant tissue stops differentiating, galls cannot be formed by a gall-maker. This explains the seasonality of leaf and bud galls, as well as the ability for stem galls arising from cambial tissue anytime during the growing season.

The Fourth Gall Law: Gall structures and locations on the plant are so species-specific that the species of the gall-maker can be identified by the gall structure alone without the need to see

A history of galls

Plant galls are remarkable plant structures that have been observed, studied and utilized since antiquity. The Greek naturalist Theophrastus (372-286 B.C.) wrote about “gall-nuts” of Syria. The Roman naturalist Pliny the Elder (23-79 A.D.) wrote about medical uses of gall extracts. In fact, much of our history was literally written in gall ink, a mixture of iron and gallic acid (3,4,5-trihydroxybenzoic acid) extracted from oak galls. The Dead Sea Scrolls were written in iron-gall ink. Leonardo da Vinci wrote his notes with it, J.S. Bach composed with it, Rembrandt and Van Gogh drew with it, and the Magna Carta was penned with gall ink. Indeed, the U.S. Declaration of Independence, the Constitution and the Bill of Rights, collectively referred to as the “Charters of Freedom” by the U.S. National Archives, were penned on parchment using iron-gall ink.

Cecidium (plural cecidia) is another name for a plant gall and is derived from the Greek *kekis*, meaning “gall nut.” Thus, cecidology translates to “the study of gall nuts” and is the study of galls and the process of gall formation; the official publication of the British Plant Gall Society is called *Cecidology*.

Cecidozoa is another name for an organism that can induce and direct gall development; it usually refers to an arthropod gall-maker. Although cecidology is a very old science, it is primarily recognized as an active branch of science by European rather than North American academia.



It is common to find the gnarled, woody, horned oak galls produced by the cynipid wasp, *Callirhytis cornigera*, affecting only a few trees in a whole row of seemingly related pin oaks (*Quercus palustris*).


the gall-maker itself. Although galls may change color and texture as gall-makers develop, and are said to be “mature” once the gall-maker completes its development, the changes are predictable.

There are more than 20 different types of “oak apple” galls, so named because of their resemblance to varying sized apples. However, only one gall wasp species, *Amphibolips quercusinanis* (syn. *A. inan-is*), produces the so-called larger empty oak-apple gall. The 1 to 1½-inch-diameter galls arise from leaf buds. The galls hold a single wasp larva and as the larva develops, the galls bear a striking resemblance to green apples with the apple-ruse made complete by reddish speckles that resemble insect damage to the “apples.” The galls are filled with soft fiber radiating spoke-like from a center kernel that houses the resident wasp larva. Once the larva pupates, the mature galls turn tannish brown and the fiber degrades, causing the galls to appear empty, thus the common name.

The Fifth Gall Law: Gall-makers are specific to certain hosts. Their activity may be confined to a plant species, or certain varieties, cultivars or provenances within a species. It is common to find the gnarled, woody, horned oak galls produced by the cynipid wasp, *Callirhytis cornigera*, affecting only a few trees in a whole row of seemingly related pin oaks (*Quercus palustris*). This observation speaks to the impact of only slight genetic differences in the gall-makers’ host. Galling activity may also be confined to a group of related species within a genus. It is uncommon for gall-makers on white oaks to also reside on red oaks. What this means for gall management is that making slight changes in tree selection may

Continued on page 27

A weevil has colonized a cedar-apple rust gall.



**We grow over
2000 different
perennials
and broker over
400 different
Clematis
varieties which
are grown by
our family as
well.**

Rijnbeek and Son is a bare-root perennial nursery based in Holland with over 75 years of experience in propagating and growing perennials.

Currently we are market leaders in Europe due to the reputation of both our finished plants and our liners.

A growing number of industry professionals in North America are getting familiar with Rijnbeek and use our bare-root plant material and provided knowledge to help them grow fantastic quality finished perennials.

New plant introductions have been done in the last years with Agapanthus, Echinacea, Gaura, Geranium, Salvia and Veronica.

We are working together with several breeders in Europe.

Most of This is also available as URC and plugs or stage 3 or 4 direct in the US

You can also find us at MANTS in Baltimore, MD, Cultivate in Columbus, OH and at all PPA meetings.

If you have any questions or would like to get more information, please feel free to contact us:

For the US Northeast: Joel Gaydos, joel@rijnbeek.com 443-326-8779

For the West Coast: Lou Vergeer, lou@rijnbeek.com 805-765-0830

For other areas: our headquarters In the Netherlands, nico@rijnbeek.com

www.rijnbeek.com





More than 20 different arthropods may wholly or partially depend upon the gnarled, woody, horned oak gall produced by the cynipid wasp, *Callirhytis cornigera*, for their livelihood.

Continued from page 25

eliminate a recurring gall problem in a landscape.

Sixth Gall Law: Insect and mite galls may house other occupants that have nothing to do with gall formation; however, these interlopers still rely on the gall for both their lodging and food. Biologists call these gall-guests “inquilines,” a term derived from the Latin *inquilinus*, which means “tenant” or “lodger.” Inquilines are divided into *cecidophages*, arthropods that feed on the gall tissue while the gall-maker is developing, and *successori*, arthropods that feed on the galls after the gall-maker moves out.

Studies have shown that more than 20 different arthropods may wholly or partially depend upon the gnarled, woody, horned oak gall produced by the cynipid wasp, *Callirhytis cornigera*, for their livelihood. Arthropods may also colonize fungal galls, such as the aforementioned fungal black knot and cedar-apple rust galls, which may introduce some confusion as to the identity of the true gall-maker! Indeed, with the wide array of possible inquilines in some galls coupled with the predators and parasitoids that feed on the gall-maker as well as the inquilines, biologists often describe galls a mini-ecosystems; it's a jungle in there!

Next up: Gall management.

Joe Boggs is an assistant professor with the Ohio State University (OSU) Extension and OSU Department of Entomology. He works as a commercial horticulture educator for OSU Extension, Hamilton County (Cincinnati). Boggs can be reached via e-mail at boggs.47@osu.edu.

Jim Chatfield is an associate professor and an extension specialist with OSU Extension in the Departments of Plant Pathology and Horticulture and Crop Science. Chatfield can be reached via e-mail at Chatfield.1@osu.edu.



There are more than 2,000 species of insect gall-makers in the U.S.

Musser Forests, Inc.

Seedlings

Transplants

Potted Liners

Container

B&B

Indiana, PA

wholesale@musserforests.com

1/800/643/8319

Fax 724/465/9893